

ASSISTED LIVING FACILITY - USAGE GUIDE



KILL CLAIMS	CONCENTRATION	DILUTION
SANITIZER For pre-cleaned, hard non-porous inanimate surfaces - Food contact surface sanitizer (No rinse) - 60 second dwell time	100 - 200 ppm	(1) 334mg tab / 1 qt (1) 3.3g tab / 2 gal (1) 13.1 tab / 10 gal
DISINFECTION & OUTBREAK CONTROL Bloodborne pathogen, Herpes & Polio, Norovirus, MRSA & Ecoli, HIV, Hep B & Hep C claims, C.diff - 4 min contact time	4306 ppm	(4) 3.3 g tab / 1 qt (4) 13.1 g tab / 1 gal
TUBERCULOSIS - 4 min contact time	5382 ppm	(5) 3.3 g tab / 1 qt

DEFINITIONS	
SANITIZING	Reducing germs on inanimate surfaces to levels considered safe by public health codes or regulations. Sanitizing may be appropriate for food service and dining areas.
DISINFECTING	Destroying or inactivating most germs on any inanimate object, but not bacterial spores. Disinfecting may be appropriate for door and cabinet handles, toilets, and other bathroom surfaces.
CONTACT TIME	The duration a surface must remain wet with a sanitizer/disinfectant to work effectively.

	MFR. NO.	DESCRIPTION	EA.
PRODUCT KIT	ESPT334MG	334MG PURTABS – Food contact surface safe sanitizer (100 tabs/box)	1 Case
	ESPT3.3G	3.3G PURTABS – Hospital grade disinfectant (200 tabs/tub)	1 Case
	ESPT13.1G	13.1G PURTABS – Hospital grade disinfectant (200 tabs/tub)	
STARTER KIT	PX200ES	PROTEXUS CORDLESS ELECTROSTATIC SPRAYER	1 Per Facility
	PX30ES	REPLACEMENT TANK	
	PX300ES	PROTEXUS ELECTROSTATIC BACKPACK SPRAYER	
	PX91ES	PX200ES Shoulder Strap	
	PX55ES	Single 60 Micron Nozzle	

*PX300ES (Backpack model) and 13.1G PURTABS may be used in conjunction with the PX200ES (Handheld model) in larger facilities.

ASSISTED LIVING FACILITY - USAGE GUIDE



ESTIMATED USAGE OF 3.3G TABS IN ASSISTED LIVING FACILITIES (ALF) - 4306PPM CONCENTRATION

SMALL ALF (0-50 BEDS)	= one plus (1+) tub of 3.3g tabs/month
MEDIUM ALF (50-100 BEDS)	= two plus (2+) tubs of 3.3g tabs/month
LARGE ALF (100+)	= three plus (3+) tubs of 3.3g tabs/month

CLEANING, SANITIZING, AND DISINFECTING FREQUENCY TABLE

AREA	CONCENTRATION	TABLET SIZE & QTY	SCHEDULE
Food Contact Surface Areas: (kitchen, dining area, cafes)	100 - 200 ppm 60 seconds	(1) 334 mg/qt	DAILY - end of day
Daily Sanitizing: public areas, salons, rehabs	100 - 200 ppm 60 seconds	(1) 334 mg/qt	DAILY: Rehab & Salon after each use
Daily Room Disinfection	1076 ppm 10 minutes	(1) 3.3 g/qt	DAILY with microfiber spray bottle
Public Area Disinfection	1076 ppm 10 minutes	(1) 3.3 g/qt	WEEKLY (daily if outbreak containment)
Terminal Clean Disinfection: After a patient is discharged, prior to a new patient entering the space, when a room is contaminated/isolated, or following renovation.	4306 ppm 4 minutes	(4) 3.3 g/qt	When room is vacated
Outbreak Control/Cold-flu Season Public Areas, containment areas, transportation buses	1076 ppm 10 minutes	(1) 3.3 g/qt	DAILY until infection is clear
Offices (staff - medical)	100 - 200 ppm 60 seconds	(1) 334 mg/qt	BI-WEEKLY OR DAILY
Transportation Buses	100 - 200 ppm 60 seconds	(1) 334 mg/qt	BI-WEEKLY OR WEEKLY
Patient Transport: Wheel Chairs & Walkers	1076 ppm 10 minutes	(1) 3.3 g/qt	AFTER EACH USE
Re-usable non critical patient care devices (blood pressure cuffs, oximeter prober)	1076 ppm 10 minutes	(1) 3.3g/qt	DAILY until infection is clear
Isolation Areas: Precaution & C.diff rooms	4306 ppm 4 minutes	(4) 3.3g/qt	After each use